



Short Communication

Early-life socioeconomic disadvantage, not current, predicts accelerated epigenetic aging of monocytes

Makeda K. Austin^a, Edith Chen^{a,b}, Kharah M. Ross^c, Lisa M. McEwen^d, Julia L. Maclsaac^d, Michael S. Kobor^{d,*}, Gregory E. Miller^{a,b,**}

^a Psychology Department, Northwestern University, United States

^b Institute for Policy Research, Northwestern University, United States

^c Psychology Department, University of California, Los Angeles, United States

^d Centre for Molecular Medicine and Therapeutics, Department of Medical Genetics, University of British Columbia, Vancouver, BC, Canada



ARTICLE INFO

Keywords:

Socioeconomic status (SES)

Epigenetic aging

Immune

ABSTRACT

Low socioeconomic status (SES) in early-life and adulthood independently contribute to increased risk for aging-related chronic diseases. One mechanistic hypothesis for these associations involves faster cellular aging of immune cells, which could plausibly contribute to chronic disease pathogenesis by compromising host resistance and/or up-regulating inflammation. However, little is known about the association between life-course SES and cellular aging. The present study examines the association of early-life and current SES with a novel biomarker of cellular aging termed the “epigenetic clock,” in monocytes. Additionally, we examine health behaviors and depressive symptoms as potential explanatory pathways. The study involved 335 participants between the ages of 15 and 55 from Vancouver, Canada and surrounding areas. Enrolled participants had to fit into four life-course SES trajectories, corresponding to low-low, low-high, high-low and high-high combinations of early-life (ages 0 to 5) and current SES respectively. Cellular aging of monocytes was measured using Horvath's DNA methylation derived measure of epigenetic age acceleration. Results indicated that socioeconomic disadvantage during early-life, but not later in life, was associated with accelerated epigenetic aging of monocytes. No early-life SES by current SES interaction was detected, suggesting that socioeconomic mobility is unrelated to epigenetic age acceleration. In path analyses, neither current health behaviors nor current depressive symptoms emerged as mediators of the early-life SES effect. These findings suggest socioeconomic disadvantage in early-life is independently predictive of cellular aging of immune cells, with potential implications for aging-related diseases later in life.

1. Introduction

Aging-related chronic diseases such as heart disease, diabetes, and cancer are among the leading causes of death worldwide. Morbidity and mortality from these conditions are disproportionately high among socioeconomically disadvantaged individuals (Braveman and Barclay, 2009). Although much evidence comes from studies of socioeconomic status (SES) during adulthood, accumulating data suggest early-life SES is also associated with age-related health problems. Independent of adulthood SES, early-life SES confers increased risk for upper respiratory infection, premature death, and morbidity and mortality from cardiovascular disease and cancer (reviewed in Miller et al., 2011). These findings suggest early-life and adulthood disadvantage increase

vulnerability to aging-related diseases through at least partially independent mechanisms.

There is considerable interest in elucidating biological mechanisms underlying socioeconomic disparities. Given the array of health problems associated with low SES, it is tempting to speculate a common mechanistic pathway. Identifying one has been challenging, because conditions that pattern by SES have diverse pathogenic features. One possibility is that disadvantage affects a general biological process, like cellular aging, and thus manifests in multiple tissues, interacting with other liabilities to increase disease risk. However, studies of telomere length, an established cellular aging metric, and life-course SES have yielded inconsistent findings (Robertson et al., 2013). Mounting evidence suggests Horvath's “epigenetic clock” is a more sensitive and

* Corresponding author at: Department of Medical Genetics, University of British Columbia, Center for Molecular Medicine and Therapeutics, BC Children's Hospital Research Institute, Canada.

** Corresponding author at: Department of Psychology, Northwestern University, 2029 Sheridan Road, Evanston, IL, 60208, United States.

E-mail addresses: msk@cmmt.ubc.ca (M.S. Kobor), greg.miller@northwestern.edu (G.E. Miller).

prognostic indicator of cellular aging than telomere length, and independently predicts mortality (for detailed comparison, see Horvath, 2013; Marioni et al., 2016). Horvath's clock aggregates DNA methylation (DNAm) at 353 age-associated CpG sites to quantify DNAm age. Epigenetic age acceleration represents variance in DNAm age not accounted for by chronological age. Positive values on this metric suggest accelerated cellular aging – i.e., that an individual is biologically older than expected – and have been associated with all-cause mortality, some cancers, and morbidity and mortality from cardiometabolic disease (Horvath, 2013; Jones et al., 2015; Marioni et al., 2016; Perna et al., 2016). One study of adults found that low SES across the life-course was associated with epigenetic age acceleration (Fiorito et al., 2017). While this was a large, multi-cohort study, it focused on older adults, a sizeable minority of whom had a history of cancer or heart disease. Here, we examined epigenetic age acceleration patterns in younger adults who were free of chronic diseases.

We hypothesized that low early-life and current SES would be associated with positive epigenetic age acceleration. We focus on early-life, based on evidence suggesting the immune system is particularly sensitive to environmental stimulation during the first years of life (Miller et al., 2011). Additionally, we considered how life-course SES trajectories, of stability or mobility, might relate to epigenetic age acceleration. We used a 2×2 design, which crossed level of disadvantage (low vs. high) with stage of life-course (early-life vs. current) and measured the epigenetic aging of participants' monocytes. We also explored potential mediating pathways. These include lifestyle characteristics – smoking, alcohol, abdominal adiposity, and inactivity – which are linked to epigenetic age acceleration (Lam et al., 2012). and depressive symptoms, a known correlate of immune dysregulation (Haroon et al., 2012).

2. Methods

Between 2009 and 2012 we enrolled 360 healthy participants from the Vancouver, British Columbia, Canada area. Recruitment and eligibility criteria are extensively detailed elsewhere (Hostinar et al., 2017). The University of British Columbia's Research Ethics Board approved this study. All adult participants provided written informed consent; for those under 18, guardians provided consent.

2.1. Design

This 2×2 design only enrolled participants who met criteria for one of four life-course SES trajectories: low early-life/low current, low early-life/high current, high early-life/low current, and high early-life/high current SES. SES was operationalized as occupational status, using the extensively validated United Kingdom's National Statistics Socioeconomic Classification (NS-SEC) (Rose et al., 2005). Early-life SES was defined as maximum parental occupational status from years 0–5. Current SES was defined as maximum household occupational status within 5 years before study enrollment. In the NS-SEC, occupational status is graded on an 8-point scale, which breaks down into three broader categories (low, middle, high). To qualify as low-SES, participants needed an NS-SEC score of > 5 , which corresponds to service and manual occupations, e.g., cleaners and transportation operatives. To qualify as high-SES, an NS-SEC score < 2 , was required, which corresponds to higher managerial and professional occupations, e.g., architects and physicians.

2.2. Measures

2.2.1. Epigenetic age acceleration

Most studies of epigenetic age acceleration focus on heterogeneous pools of cells, creating uncertainty about which leukocyte sub-population(s) are involved. To circumvent this problem we focused on monocytes, a cell population sensitive to life-course SES and implicated

in multiple age-related diseases (Miller et al., 2011). Following an overnight fast, antecubital blood was collected into Cell Preparation Tubes (BD Biosciences). Peripheral blood mononuclear cells (PBMC) were isolated by density-gradient centrifugation. From the resulting pellet, monocytes were isolated by immune-magnetic capture using antibodies against CD14 (Miltenyi AutoMACS). Monocytes were lysed, homogenized, and stored at -80°C . Next, genomic DNA was isolated using a column-based method and assessed for quality and quantity using a NanoDrop. Approximately 750 ng of genomic DNA was used for bisulfite treatment for sequence-based differentiation of methylated cytosine nucleotides. Bisulfite converted DNA samples were randomized and processed using the Illumina 450 K array. Horvath's clock was estimated with a publicly available R script. Epigenetic age acceleration was calculated by extracting residuals from a linear model wherein DNAm age was regressed onto chronological age (Horvath, 2013).

2.2.2. Potential mediators

Past month depressive symptoms were assessed with the Center for Epidemiologic studies Depression scale (CES-D) brief version (Andresen, 1994). (Cronbach's $\alpha = 0.84$). Participants reported daily number of cigarettes smoked and weekly alcoholic drinks, coded as none (0), < 10 (1), or > 10 (2). Physical activity was measured as weekly minutes of brisk activity (Paffenbarger et al., 1993). Abdominal adiposity was operationalized as continuous waist circumference. Participants self-reported their age, sex, and race/ethnicity.

2.3. Data preparation

Variables were examined for outliers before analyses. Values exceeding four mean standard deviations were Winsorized with the value at the 99.9th percentile (waist circumference: $n = 3$). Physical activity was skewed and normalized with a square-root transformation.

2.4. Missing data

Of 360 participants, 25 (6.9%) were missing data for epigenetic aging due to technical problems with venipuncture, sample processing, or assay procedures. Between 0.3% and 4.5% of participants were missing data for smoking: $n = 15$, alcohol use: $n = 10$, physical activity: $n = 10$, waist circumference: $n = 1$, and depressive symptoms: $n = 8$. Therefore, analytic N s ranged from 320 to 335.

3. Results

Table 1 provides characteristics of the sample. Consistent with previous studies, chronological age and DNAm age were strongly correlated ($r(333) = 0.93$, $p = 1.1 \times 10^{-144}$). As expected, the derived epigenetic age acceleration metric was uncorrelated with chronological age, $r(333) = 0.00$, $p = 0.996$.

Next, we used ANOVAs to address whether life-course SES was associated with epigenetic age acceleration. Crude models contained variables reflecting early-life SES, current SES, and their interaction; adjusted ANCOVA models also included the covariates age, race, and sex. Hypotheses about low SES were partially supported. We observed a main effect of early-life SES in both unadjusted, $F(1, 334) = 4.50$, $p = .035$, $d = .24$, and adjusted models $F(1, 334) = 4.33$, $p = .038$, $d = .23$. Participants with low early-life SES showed more epigenetic age acceleration compared to those with high early-life SES (Fig. 1). The difference corresponded to an epigenetic age acceleration of 0.93 years. In contrast, there was no main effect of current SES in unadjusted, $F(1, 333) = 0.10$, $p = .748$, $d = 0.03$ or adjusted models, $F(1, 333) = 0.07$, $p = 0.790$, $d = 0.03$. There also was no significant interaction between early-life and current SES in either unadjusted, $F(1, 333) = 0.17$, $p = 0.693$, $d = 0.06$, or adjusted models, $F(1, 333) = 0.26$, $p = 0.61$, $d = 0.06$.

Table 1
Characteristics of the sample (N = 335).

	Life-course SES Trajectory				Main effect of early-life SES		Main effect of current SES	
	Low/Low n = 96	Low/High n = 85	High/Low n = 67	High/High n = 87	Estimate	p	Estimate	p
Age (years)								
15 – 24 (%)	19.8	2.4	19.4	18.4	F = 6.45	.012	F = 3.59	.059
25 – 39 (%)	42.7	44.7	47.8	44.8				
40 – 55 (%)	37.5	52.9	32.8	36.8				
Female (%)	50	58.3	58.2	54	X ² = 0.17	.680	X ² = 0.20	.654
White (%)	67.7	66.7	80.6	80.5	X ² = 7.34	.007	X ² = 0.03	.863
High abdominal adiposity (%)	13.0	15.0	14.7	14.7	F = 0.40	.526	F = 0.29	.589
Heavy smoker (%)	17.7	2.5	16.9	3.5	F = 0.18	.668	F = 20.64	< .001
Heavy drinker (%)	10.9	11.1	18.5	16.1	F = 1.36	.244	F = 0.06	.802
Physical activity (min/week)	165.5	161.0	196.4	181.5	F = 2.30	.131	F = 0.28	.598
	(204.3)	(142.1)	(180.4)	(174.1)				
Current Depressive Symptoms	9.2	7.0	9.6	6.6	F < 0.01	.977	F = 19.50	< .001
	(5.7)	(4.8)	(6.24)	(4.53)				

Note: Participants were recruited to fit into 1 of 4 distinct life-course SES trajectories; low early-life /low current (Low/Low), low early-life /high current (Low/High), high early-life /low current (High/Low), and high early-life /high current (High/High). Mean (SD) reported unless otherwise noted. F statistic and p-value reported for 2-way ANOVA analyses examining the main effects of early SES and current SES and their interaction. Chi-square analysis was used in the case of binary IVs and DVs. High risk abdominal adiposity is categorized as > 102 cm for men and > 88 cm for women based upon NIH risk classification for type 2 diabetes, hypertension, and cardiovascular disease (https://www.nhlbi.nih.gov/health/educational/lose_wt/BMI/bmi_dis.htm). Heavy smoker was defined as smoking ≥ 10 cigarettes per day. Heavy drinker was defined as consuming ≥ 10 drinks per week. Current depressive symptoms were assessed with the Center for Epidemiologic Studies Depression Scale (CES-D).

Our sample included 7 participants under 18, and 50 participants under 25 who may have not yet achieved their maximum adulthood SES. These participants could have diluted any current SES effects. To examine this, we re-conducted adjusted analyses excluding all 50 below age 25. The results were unchanged: even in this older sample, no main effect of current SES ($F(1, 333) = 0.01, p = .917, d = 0.01$) or early-life x current SES interaction ($F(1, 333) = 0.06, p = .803, d = 0.03$) was apparent.

Next, we examined lifestyle characteristics and depressive symptoms as potential mediating pathways between early-life SES and epigenetic age acceleration. These analyses were performed with the PROCESS macro, which uses regression-based path-analysis to estimate direct and indirect pathways (Hayes, 2013), and bootstrap methods to obtain bias-corrected indirect effects. Models controlled for race, sex, and age.

For all indirect pathways considered, the 95% confidence intervals (CI) around the estimated coefficient included 0, inconsistent with mediation. This was observed in unadjusted and adjusted analyses for smoking $b = -0.001, CI (-0.031, 0.032)$, alcohol use $b = 0.001, CI (-0.011, .030)$ physical activity $b = -0.002, CI (-0.034, 0.007)$ waist

circumference $b = 0.001, CI (-0.011, 0.030)$, and depressive symptoms $b = -0.001, CI (-0.022, 0.012)$.

4. Discussion

We observed that low early-life SES, but not current SES, was associated with accelerated epigenetic aging of monocytes. Consistent with previous research (Fiorito et al., 2017), we did not find that upward socioeconomic mobility across the life-course substantially attenuated this association. These findings contribute to a growing literature highlighting the association between early-life SES and lifelong health. These patterns suggest the possibility that early-life experiences get “embedded” in the monocyte epigenome in a durable manner. However, the embedding scenario cannot be evaluated here with certainty, due to our focus on monocytes, which are non-dividing cells with short life cycles (Miller et al., 2011). A next step is to investigate whether these patterns are evident in long-living cells such as hematopoietic stem cells, and other monocyte lineage precursors.

The effect size for the association between low early-life SES and epigenetic age acceleration was relatively small, amounting to 0.93




Fig. 1. Epigenetic age acceleration, calculated by extracting residuals from a linear model of DNA methylation age onto chronological age, varied by life-course trajectories of early-life and current socioeconomic status. Low early-life/low current (Low/Low), low early-life/high current (Low/High), high early-life/low current (High/Low), high early-life/high current (High/High).

years. Comparable studies suggest one year of accelerated epigenetic aging corresponds to a 4.4% increased risk of all-cause mortality (Perna et al., 2016).

We did not observe differences in epigenetic age acceleration by current SES. Although this result may surprise readers, it converges with a meta-analysis which found no cumulative evidence of an association between current SES and telomere length, and with findings suggesting that PBMC DNAm was associated with childhood but not adulthood SES (Lam et al., 2012). Nevertheless, it is possible that characteristics of our sample, such as the wide age range, obscured genuine differences.

Lastly, we examined mediators that might explain the link between early-life SES and epigenetic age acceleration. There was no evidence that lifestyle factors or depressive symptoms played such roles, suggesting early-life SES influences cellular aging through other pathways. However, we only assessed these processes later in life, and they may have played mediating roles in earlier years. Prospective studies that originate in childhood, and examine other plausible mediators, such as environmental pathogen exposure and childhood maltreatment, might be useful in linking specific childhood experiences to epigenetic age acceleration (Braveman and Barclay, 2009; Miller et al., 2011).

Due to our cross-sectional design and retrospective assessment, we cannot infer causal relationships. Future studies should expand on our findings with more population representative samples and include additional metrics of cellular aging, e.g., telomere length. Given our SES group study design, we were unable to examine a continuous linear approach. Future studies should examine this and potential differential influences of maternal vs. paternal SES. Lastly future studies should explore how SES interacts with other domains of social stratification that confer overlapping life-course disadvantage. For instance, among Whites, upward socioeconomic mobility is associated with better health in adulthood. However, among racial/ethnic minority groups from disadvantaged backgrounds, mobility forecasts worse health (Gaydos et al., 2017).

5. Conclusion

This study provides evidence that socioeconomic disadvantage in early-life is associated with accelerated epigenetic aging of monocytes. This suggests socioeconomic disadvantage in the first years of life may have lasting implications for cellular aging, a potentially common pathological mechanism underlying socioeconomic disparities across aging-related diseases.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgements

We thank the study participants and the NIH Grants R01 HD058502 and R01 HL122328 for supporting this project. We also thank Rachel Edgar and Nicole Gladish for their assistance on this project.

References

- Andresen, E.M., 1994. Screening for depression in well older adults: evaluation of a short form of the CES-D. *Am. J. Prevent. Med.* 10 (2), 77–84.
- Braveman, P., Barclay, C., 2009. Health disparities beginning in childhood: a life-course perspective. *Pediatrics* 124 (Suppl. 3), S163–S175. <https://doi.org/10.1542/peds.2009-1100D>.
- Fiorito, G., Polidoro, S., Dugué, P., Kivimaki, M., Ponzi, E., Matullo, G., Simonetta, G., Assumma, M.B., Georgiadis, P., Kyrtopoulos, S.A., Krogh, V., Palli, D., Panico, S., Sacerdote, C., Tumino, R., Chadeau-Hyam, M., Stringhini, S., Severi, G., Hodge, A.M., Giles, G.G., Marioni, R., Linner, R.K., O'Halloran, A.M., Kenny, R.A., Layte, R., Baglietto, L., Robinson, O., McCrory, C., Milne, R.L., Vineis, P., 2017. Social adversity and epigenetic aging: a multi-cohort study on socioeconomic differences in peripheral blood DNA methylation. *Sci. Rep.* 7 (1). <https://doi.org/10.1038/s41598-017-16391-5>.
- Gaydos, L., Schorpp, K., Chen, E., Miller, G.E., Harris, K.M., 2017. College completion predicts lower depression but higher metabolic syndrome among disadvantaged minorities in young adulthood. *Proc. Natl. Acad. Sci. U. S. A.* 115 (1), 109–114. <https://doi.org/10.1073/pnas.1714616114>.
- Haron, E., Raison, C.L., Miller, A.H., 2012. Psychoneuroimmunology meets neuropsychopharmacology: translational implications of the impact of inflammation on behavior. *Neuropsychopharmacology* 37 (1), 137–162. <https://doi.org/10.1038/npp.2011.205>.
- Hayes, A.F., 2013. *Introduction to Mediation, Moderation, and Conditional Process Analysis*. The Guilford Press, New York.
- Horvath, S., 2013. DNA methylation age of human tissues and cell types. *Genome Biol.* 14 (10), R115. <https://doi.org/10.1186/gb-2013-14-10-r115>.
- Hostinar, C.E., Ross, K.M., Chen, E., Miller, G.E., 2017. Early-life socioeconomic disadvantage and metabolic health disparities. *Psychosom. Med.* 79 (5), 514–523. <https://doi.org/10.1097/psy.0000000000000455>.
- Jones, M.J., Goodman, S.J., Kobor, M.S., 2015. DNA methylation and healthy human aging. *Aging Cell* 14 (6), 924–932. <https://doi.org/10.1111/ace1.12349>.
- Lam, L.L., Emberly, E., Fraser, H.B., Neumann, S.M., Chen, E., Miller, G.E., Kobor, M.S., 2012. Factors underlying variable DNA methylation in a human community cohort. *Proceedings of the National Academy of Sciences of the United States of America* 109 (Suppl. 2), 17253–17260. <https://doi.org/10.1073/pnas.1121249109>.
- Marioni, R.E., Harris, S.E., Shah, S., Mcrae, A.F., Zglinicki, T.V., Martin-Ruiz, C., Wray, N.R., Visscher, P.M., Deary, I.J., 2016. The epigenetic clock and telomere length are independently associated with chronological age and mortality. *Int. J. Epidemiol.* 45 (2), 424–432. <https://doi.org/10.1093/ije/dyw041>.
- Miller, G.E., Chen, E., Parker, K.J., 2011. Psychological stress in childhood and susceptibility to the chronic diseases of aging: moving towards a model of behavioral and biological mechanisms. *Psychol. Bull.* 137 (6), 959–997. <https://doi.org/10.1037/a0024768>.
- Paffenbarger, R.S., Blair, S.N., Lee, I.M., Hyde, R.T., 1993. Measurement of physical activity to assess health effects in free-living populations. *Med. Sci. Sports Exerc.* 25, 60–70.
- Perna, L., Zhang, Y., Mons, U., Hollecsek, B., Saum, K.-U., Brenner, H., 2016. Epigenetic age acceleration predicts cancer, cardiovascular, and all-cause mortality in a German case cohort. *Clin. Epigenet.* 8, 64. <https://doi.org/10.1186/s13148-016-0228-z>.
- Robertson, T., Batty, G.D., Der, G., Fenton, C., Shiels, P.G., Benzeval, M., 2013. Is socioeconomic Status associated with biological aging as measured by telomere length? *Epidemiol. Rev.* 35 (1), 98–111. <https://doi.org/10.1093/epirev/mxs001>.
- Rose, D., Pevalin, D.J., O'Reilly, K., 2005. *The National Statistics Socioeconomic Classification: Origins, Development and Use*. Palgrave Macmillan, Basingstoke.